

MODELING FLOOD HAZARDS IN AMBON CITY WATERSHEDS: CASE STUDIES OF WAI BATU GANTUNG, WAI BATU GAJAH, WAI TOMU, WAI BATU MERAH AND WAI RUHU

Heinrich Rakuasa^{1*}, Philia Christi Latue²

¹National Research Tomsk State University, Russian Federation

²Universitas Pattimura, Indonesia

heinrichrakuasa01@gmail.com¹

Received: 10-04-2024

Revised: 23-04-2024

Approved: 30-04-2024

ABSTRACT

Flood hazard modeling in watersheds is an important step in natural disaster risk mitigation, especially in vulnerable areas such as Ambon City. This research focused on the Wai Batu Gantung, Wai Batu Gajah, Wai Tomu, Wai Batu Merah, and Wai Ruhu watersheds, using JRC Global Surface Water Mapping Layers data, NASA SRTM Digital Elevation 30 m data, and USGS Landsat 8 Level 2, Collection 2, Tier 1 data analyzed on the Google Earth Engine (GEE) platform. Prediction of built-up land in flood-prone areas was conducted by utilizing flood history analysis, hydrological modeling, and flood zone mapping. The results show that flood hazard modeling provides a better understanding of flood risk, assists in the development of safer land use planning, and increases public awareness of flood risk in Ambon City. It is hoped that the results of this research can contribute to flood risk management and sustainable regional development in the future.

Keywords: Ambon, Flood, Geogle Eart Engine, Watershed

INTRODUCTION

Flood hazard modeling in river basins is an important step in natural disaster risk mitigation, especially in vulnerable areas such as Ambon City (Rakuasa et al., 2022; Muin & Rakuasa 2023). Ambon City, located on Ambon Island, Maluku, is one of the important cities in Eastern Indonesia that often experiences floods due to natural and anthropogenic factors (Rakuasa et al., 2022). By using advanced technology such as Google Earth Engine, researchers can conduct accurate and detailed modeling of flood-prone areas around major rivers in Ambon City (Sugandhi & Rakuasa 2023; Manakane et al., 2023). This case study focuses on several major rivers in Ambon City, namely Wai Batu Gantung, Wai Batu Gajah, Wai Tomu, Wai Batu Merah, and Wai Ruhu. These rivers play a vital role in the lives of local communities, but they also carry the potential for serious flooding hazards, especially during the rainy season (Rakuasa et al., 2022). Therefore, modeling flood hazards along these rivers is essential to assist the government and communities in mitigating and managing disaster risks (Rakuasa & Latue 2023; Latue et al., 2023)

Google Earth Engine is a platform that enables sophisticated spatial analysis and massive processing of satellite images (Letedara et al., 2023). By utilizing this technology, researchers can access high-resolution satellite imagery data from multiple sources to map and analyze topographic conditions, land use, vegetation, and other factors that affect water flow along the rivers being studied (Cui et al. 2023). Flood hazard modeling with Google Earth Engine allows researchers to estimate potential flood vulnerabilities along these rivers in a high level of detail. Through careful analysis, they were able to identify flood hotspots, determine water flow patterns, and predict

the potential impact of flooding on surrounding settlements and infrastructure.

The modeling method used in this study involves processing and analyzing satellite image data using specialized algorithms available in the Google Earth Engine. The steps include image segmentation, extraction of important features, such as rivers, and mapping of water flow patterns using a proven hydrological model. The results of this flood hazard modeling are expected to provide a better understanding of the water flow dynamics and potential flood risks in the studied watershed. This information can be used by local governments in spatial planning, disaster risk management, and decision-making related to infrastructure and settlement development (Gustavo and Rakuasa 2023; Septory et al., 2023). In addition, the results of this modeling can also be the basis for the development of a more effective and accurate flood early warning system. By utilizing real-time satellite imagery data and calibrated hydrological models, early warning systems can provide early warnings to the public so that they can take preventive or evacuation measures in a timely manner (Rakuasa et al., 2023).

However, flood hazard modeling also has several challenges and limitations. One of them is the limitation of input data, especially in terms of satellite image resolution and topographic data accuracy (Halder and Bose 2024). In addition, the hydrological models used also need to be calibrated and validated with field data to ensure their accuracy in predicting flood risks (Zhao et al. 2021). Nonetheless, the use of Google Earth Engine in flood hazard modeling offers great potential in improving the understanding and mitigation of disaster risk in Ambon City and other areas prone to flooding (Papaioannou et al. 2016). By continuing to develop and improve modeling methods and increasing access to quality satellite imagery data, it is hoped that flood risk mitigation and management efforts can become more effective and efficient in the future.

RESEARCH METHODS

This research was conducted in the watershed of Wai Batu Gantung, Wai Batu Gajah, Wai Tomu, Wai Batu Merah and Wai Ruhu, Ambon City. This research uses JRC Global Surface Water Mapping Layers, v1.4, NASA SRTM Digital Elevation 30 m data, and USGS Landsat 8 Level 2, Collection 2, Tier 1 data analyzed on the Google Earth Engine (GEE) platform. Google Earth Engine is a cloud computing platform developed by Google for large-scale geospatial data analysis and processing (Hamidi et al. 2023). By providing access to a wide range of satellite image datasets and powerful geospatial analysis tools, Google Earth Engine enables users to perform modeling, mapping, and monitoring of the earth's environment with high efficiency (Mai Sy et al. 2023). The platform facilitates temporal and spatial analysis, predictive model building, as well as interactive visualization of geospatial data, making it a very useful tool for researchers, scientists, and practitioners in various fields to run complex geospatial analyses without the need to download or process data locally.

The variables used in this study consisted of distance from the river, Normalized Difference Water Index (NDWI), elevation, Topographic Position Index (TPI) and Normalized Difference Vegetation Index (NDVI), all of which were analyzed on the Google Earth Engine (GEE) platform; <https://earthengine.google.com>. JRC Global Surface Water Mapping Layers data in Google Earth Engine (GEE) can produce Normalized Difference Water Index (NDWI) data which is very important for flood analysis [19]. NDWI is an index used to monitor and identify water surfaces, including

ivers, lakes and other bodies of water [20]. By utilizing NDWI data from JRC, researchers and practitioners can accurately and quickly identify waterlogged areas during flood events. This information is essential for flood monitoring, mapping and risk management, enabling relevant parties to respond quickly and effectively in emergency situations and plan appropriate mitigation measures to reduce the impact of future floods (Baig et al. 2013)

NASA SRTM Digital Elevation 30m data and USGS Landsat 8 Level 2 data are used to generate elevation and TPI (Topographic Position Index) which are important in flood analysis. Elevation data from SRTM allows for accurate topographic modeling, while Landsat 8 Level 2 data can be used to extract information on land texture and structure that affect surface water flow (Kelleher and McPhillips 2020). By utilizing both, users can identify low areas susceptible to inundation during floods and determine the relative topography of an area to its surroundings through TPI, thus providing more comprehensive insights in anticipating and managing flood risks (Sivanpillai et al. 2021).




Figure 1. Workflow

USGS Landsat 8 Level 2, Collection 2, Tier 1 data in Google Earth Engine (GEE) provides Normalized Difference Vegetation Index (NDVI) data that is critical for flood analysis. NDVI is an index used to measure the amount and quality of vegetation in an area based on the reflectance of near infrared and red light from the ground surface. Using NDVI data from Landsat 8, researchers and practitioners can evaluate the state of vegetation before and after flooding, which provides valuable insights into the impact of flooding on ecosystems (Soltani et al. 2021). This information can be used to evaluate environmental damage, plan habitat restoration, and identify areas that may experience soil erosion or flooding after a flood event, enabling more precise and effective mitigation and adaptation effort (Soltani et al. 2021). The flood analysis process in

Google Earth Engine can be seen in Figure 1.

These variables are considered important in this study because, distance from the river is relevant as areas closer to the river tend to be more vulnerable to flooding. NDWI provides a clear visual indication of waterlogged areas, while elevation enables water flow modeling and identification of low areas prone to inundation (Aires et al. 2020). TPI provides an understanding of the relative topography of an area, while NDVI provides information on vegetation that can affect drainage and water absorption. The use of GEE allows for efficient processing and analysis of these data, enabling faster and more informed decision-making in flood risk mitigation and adaptation. All variables were then given weights and values before being overlaid on the Google Earth Engine (GEE) platform to produce flood-prone areas and then overlaid with data on the distribution of built-up land in the Wai Ruhu watershed to determine settlements that are predicted to be affected by flooding in the Wai Ruhu watershed, Ambon City. The whole process of analyzing flood-prone areas was carried out on the Google Earth Engine (GEE) Platform with a modified script based on previous research (Aires et al. 2020).

RESULTS AND DISCUSSION

Flood Hazard Level

Flood hazard level is a measure or classification used to assess how much threat or potential damage flooding can cause in a given area. The flood hazard level usually includes several factors, including rainfall intensity, water flow velocity, topography of the area, infrastructure condition, and other human factors. This helps governments, environmental experts, and communities to take appropriate prevention and countermeasures to reduce flood risks and protect lives and property. Factors such as extreme rainfall, topography, land use, drainage systems, and river conditions can affect the level of flood vulnerability of an area. The results of the analysis of the flood hazard level of the Wai Batu Gantung, Wai Batu Gajah, Wai Tomu, Wai Batu Merah and Wai Ruhu watersheds on the Google Earth Engine (GEE) platform show that the area with low hazard level is 2,662.65 ha, medium hazard is 1,739.60 ha and high hazard is 543.79 ha. The map of flood hazard levels in the watersheds of Wai Batu Gantung, Wai Batu Gajah, Wai Tomu, Wai Batu Merah and Wai Ruhu can be seen in Figure 2.



Figure 2. Flood Hazard Map

Areas with high flood hazard levels are marked in red, medium flood hazard with medium color, low flood hazard marked in green. The flood hazard level of Wai Batu Gantung watershed consists of low hazard of 63.21 ha, medium hazard of 142.36 ha and high hazard of 73.25 ha. The flood hazard level of Wai Wai Ruhu watershed consists of low hazard of 278.44 ha, medium hazard of 278.44 ha and high hazard of 87.74 ha. The flood hazard level of Wai Wai Batumerah watershed consists of low hazard of 75.03 ha, medium hazard of 170.22 ha and high hazard of 38.20 ha. The flood hazard level of Wai Tomu watershed consists of low hazard of 114.03 ha, medium hazard of 91.56 ha and high hazard of 80.58 ha. The flood hazard level of Wai Batu Gajah watershed consists of low hazard of 65.39 ha, medium hazard of 61.22 ha and high hazard of 78.39 ha. Assessing the level of flood hazard is important to assist in flood risk mitigation planning, developing early warning systems, and increasing preparedness in the face of flood disasters. By understanding the level of flood vulnerability, relevant parties can identify vulnerable areas and take preventive or adaptive measures to reduce the adverse impacts of flooding.

Predicted built-up land located in flood hazard areas

Predictions of potential built-up land in flood-prone areas of the Wai Batu Gantung, Wai Batu Gajah, Wai Tomu, Wai Batu Merah, and Wai Ruhu watersheds can be made using historical flood analysis, hydrological modeling, and flood zone mapping. By combining historical flood data, geographical characteristics of the rivers and topography in the area, and predictions of potential water flows in flood scenarios, experts can identify areas most vulnerable to flooding, assisting stakeholders in safer and more effective land use planning in the future. Predicting flood-affected residential areas is the process of estimating areas around rivers or other flood-prone areas that are likely to be inundated or affected during a flood. The results of the analysis of the flood vulnerability level of the Wai Batu Gantung, Wai Batu Gajah, Wai Tomu, Wai Batu Merah, and Wai Ruhu watersheds on the Google Earth Engine (GEE) platform are then overlaid with built-up land data extracted from Normalized Difference Vegetation Index (NDVI) data which shows that residential areas are in flood-prone areas. The spatial map of flood-affected settlements can be seen in Figure 3.




Figure 3. Map of Flood Affected Settlements

The analysis results show that settlements predicted to be in high flood hazard areas are 543.79 ha, settlements predicted to be in moderate hazard areas are 1,739.60 ha and settlements predicted to be in low flood hazard areas are 2,662.65 ha. Settlements predicted to be affected by flood hazards in the Wai Wai Ruhu watershed consist of low hazard of 1048.45 ha, medium hazard of 725.23 ha and high hazard of 165.34 ha. Settlements predicted to be affected by flood hazards in the Wai Batu Gantung watershed consist of low hazard of 538.98 ha, medium hazard of 425.25 ha and high hazard of 125.63 ha. Settlements predicted to be affected by flood hazards in the Wai Tomu watershed consist of low hazard of 406.70 ha, medium hazard of 126.13 ha and high hazard of 87.79 ha. Settlements predicted to be affected by flood hazards in the Wai Batu Gajah watershed consist of a low hazard of 326.26 ha, a medium hazard of 205.74 ha and a high hazard of 109.52 ha. Settlements predicted to be affected by flood hazards in the Wai Batumerah watershed consist of a low hazard of 342.26 ha, a medium hazard of 257.25 ha and a high hazard of 55.52 ha.

The flood hazard modeling study in the Ambon watershed, particularly in the case of Wai Batu Gantung, Wai Batu Gajah, Wai Tomu, Wai Batu Merah, and Wai Ruhu, provided several significant benefits: (1) Better Understanding of Flood Risk: The study helps in understanding the potential flood risk in the area by analyzing the pattern and intensity of flooding that has occurred and predicting the potential for future flooding (Muin & Rakuasa 2023). This enables the government and stakeholders to take appropriate mitigation measures (Muin & Rakuasa 2023; Rakuasa & Rinaldi 2023). Development of Better Land Use Planning: By understanding the zones most vulnerable to flooding, the modeling can assist in the development of safer and more sustainable land-use planning (Muin et al., 2023). This includes restrictions on development in high-risk areas as well as the promotion of better drainage infrastructure (Sugandhi et al., 2023). Preparation for Climate Change: By considering climate change in the modeling, the study helps in preparing for the impacts of floods that may become more frequent or intense in the future due to changes in extreme weather (Wulandari et al., 2023; Latue & Rakuasa 2022). Public Awareness Raising: Information from this study can be used to raise public awareness about flood risks in the area (Rakuasa and Mehdila 2023). This can help people to take personal measures in preparation for flooding, such as the development of evacuation plans and investment in self-protection infrastructure (Rakuasa and Mehdila 2023). Thus, the flood modeling study in the Ambon watershed provides a solid foundation for better decision-making in flood risk management, environmental protection, and sustainable regional development.

CONCLUSION

It can be concluded that flood hazard modeling provides significant benefits in understanding flood risks, developing safer land use planning, preparing for climate change, and raising public awareness. By analyzing existing flood patterns and predicting potential future floods, this research provides a solid basis for appropriate mitigation measures, sustainable infrastructure development, and improved flood preparedness. It is hoped that the results of this research can make a positive contribution to flood risk management, environmental protection and sustainable regional development in the future.

REFERENCES

- Aires, Uilson Ricardo Venâncio et al. 2020. 'The Use of the Normalized Difference Vegetation Index to Analyze the Influence of Vegetation Cover Changes on the Streamflow in the Manhuaçu River Basin, Brazil'. *Water Resources Management* 34(6): 1933–49. <https://link.springer.com/10.1007/s11269-020-02536-1>.
- Baig, Muhammad Hasan Ali et al. 2013. 'Comparison of MNDWI and DFI for Water Mapping in Flooding Season'. In *2013 IEEE International Geoscience and Remote Sensing Symposium - IGARSS*, IEEE, 2876–79. <http://ieeexplore.ieee.org/document/6723425/>.
- Cui, Jiaqi et al. 2023. 'Extraction of Information on the Flooding Extent of Agricultural Land in Henan Province Based on Multi-Source Remote Sensing Images and Google Earth Engine'. *Agronomy* 13(2): 355. <https://www.mdpi.com/2073-4395/13/2/355>.
- Gustavo, Geoffrey John Pascal, and Heinrich Rakuasa. 2023. 'Disaster Education and the Role of Geographers: A Step Toward a Disaster Resilient Ambon City: A Review'. *Journal of Education Method and Learning Strategy* 1(03): 183–92. <https://risetpress.com/index.php/jemls/article/view/238>.
- Halder, Subhra, and Suddhasil Bose. 2024. 'Sustainable Flood Hazard Mapping with GLOF: A Google Earth Engine Approach'. *Natural Hazards Research*. <https://linkinghub.elsevier.com/retrieve/pii/S2666592124000027>.
- Hamidi, Ebrahim et al. 2023. 'Fast Flood Extent Monitoring With SAR Change Detection Using Google Earth Engine'. *IEEE Transactions on Geoscience and Remote Sensing* 61: 1–19. <https://ieeexplore.ieee.org/document/10026287/>.
- Heinrich Rakuasa, Daniel A Sihasale, Marhelin C Mehdila, Anelia P Wlary. 2022. 'Analisis Spasial Tingkat Kerawanan Banjir Di Kecamatan Teluk Ambon Baguala, Kota Ambon'. *Jurnal Geosains dan Remote Sensing (JGRS)* 3(2): 60–69.
- Kelleher, Christa, and Lauren McPhillips. 2020. 'Exploring the Application of Topographic Indices in Urban Areas as Indicators of Pluvial Flooding Locations'. *Hydrological Processes* 34(3): 780–94. <https://onlinelibrary.wiley.com/doi/10.1002/hyp.13628>.
- Latue, P. C., & Rakuasa, H. 2032. 'Identification of Flood-Prone Areas Using the Topographic Wetness Index Method in Fena Leisela District, Buru Regency'. *Journal Basic Science and Technology* 12(2).
- Latue, P. C., Imanuel Septory, J. S., Somae, G., & Rakuasa, H. 2023. 'Pemodelan Daerah Rawan Banjir Di Kecamatan Sirimau Menggunakan Metode Multi-Criteria Analysis (MCA)'. *Jurnal Perencanaan Wilayah Dan Kota* 18(1): 10–17.
- Letedara, R., Manakane, S. E., Latue, P. C., & Rakuasa, H. 2023. 'Analisis Spasio-Temporal Perubahan Suhu Permukaan Daratan Pulau Letti Tahun 2013 Dan 2023 Menggunakan Data Citra Ladsat 8 OLI/TIRS Pada Google Earth Engine'. *Larisa Penelitian Multidisiplin* 1(1): 36–42.
- Mai Sy, Hung et al. 2023. 'Urban Flood Risk Assessment Using Sentinel-1 on the Google Earth Engine: A Case Study in Thai Nguyen City, Vietnam'. *Remote Sensing Applications: Society and Environment* 31: 100987. <https://linkinghub.elsevier.com/retrieve/pii/S2352938523000691>.
- Manakane, S. E., Latue, P. C., Somae, G., & Rakuasa, H. 2023. 'Flood Risk Modeling in Buru Island, Maluku Province, Indonesia Using Google Earth Engine: Pemodelan Risiko Banjir Di Pulau Buru, Provinsi Maluku, Indonesia Dengan Menggunakan

- Mesin Google Earth'. *MULTIPLE: Journal of Global and Multidisciplinary* 1(2): 80–87.
- Muin, A., & Rakuasa, H. 2023a. 'Evaluasi Rencana Tata Ruang Wilayah Kota Ambon Berdasarkan Aspek Kerawanan Banjir'. *ULIL ALBAB: Jurnal Ilmiah Multidisiplin* 2(5): 1727–1738.
- . 2023b. 'Pemetaan Daerah Rawan Banjir Di Desa Lokki Kecamatan Huamual Kabupaten Seram Bagian Barat.' *Gudang Jurnal Multidisiplin Ilmu* 1(2): 47–52.
- Muin, A., & Rakuasa, H. 2023c. 'Pemanfaat Geographic Artificial Intelligence (Geo-AI) Untuk Identifikasi Daerah Rawan Banjir Di Kota Ambon'. *Gudang Jurnal Multidisiplin Ilmu* 1(2): 58-63.
- Muin, A., Somae, G., & Rakuasa, H. 2023. 'Analisis Potensi Genangan Banjir Di Kecamatan Siwalalat, Kabupaten Seram Bagian Timur Berdasarkan Topographic Wetness Index'. *ULIL ALBAB: Jurnal Ilmiah Multidisiplin* 2(5): 1800–1806.
- Papaioannou, G., A. Loukas, L. Vasiliades, and G. T. Aronica. 2016. 'Flood Inundation Mapping Sensitivity to Riverine Spatial Resolution and Modelling Approach'. *Natural Hazards* 83(S1): 117–32. <http://link.springer.com/10.1007/s11069-016-2382-1>.
- Rakuasa, H., & Rinaldi, M. 2023. 'Pemetaan Daerah Potensi Genangan Banjir Di Kecamatan Gunugputri, Kabupaten Bogor Menggunakan Data DEMNAS Dengan Metode Topographic Wetness Index'. *ULIL ALBAB: Jurnal Ilmiah Multidisiplin* 2(6): 2269-2280.
- Rakuasa, H., Helwend, J. K., & Sihasale, D. A. 2022. 'Pemetaan Daerah Rawan Banjir Di Kota Ambon Menggunakan Sistem Informasi Geografis'. *Jurnal Geografi: Media Informasi Pengembangan dan Profesi Kegeografian* 19(2): 73–82.
- Rakuasa, H., Somae, G., & Latue, P. C. 2023. 'Pemetaan Daerah Rawan Banjir Di Desa Batumerah Kecamatan Sirimau Kota Ambon Menggunakan Sistem Informasi Geografis'. *ULIL ALBAB: Jurnal Ilmiah Multidisiplin* 2(4): 1642–53.
- Rakuasa, Heinrich, and Philia Christi Latue. 2023. 'ANALISIS SPASIAL DAERAH RAWAN BANJIR DI DAS WAE HERU, KOTA AMBON'. *Jurnal Tanah dan Sumberdaya Lahan* 10(1): 75–82. <https://jtsl.ub.ac.id/index.php/jtsl/article/view/845>.
- Rakuasa, Heinrich, and Marhelin Ch Mehdila. 2023. 'Penerapan Pendidikan Mitigasi Bencana Gempa Bumi Untuk Siswa Dan Guru Di SD Negeri 1 Poka, Kota Ambon, Provinsi Maluku'. *Jurnal Pengabdian Masyarakat Indonesia* 3(3): 441–46. <https://jpmi.journals.id/index.php/jpmi/article/view/1138>.
- Rakuasa, Heinrich, Daniel Anthoni Sihasale, and Philia Christi Latue. 2022. 'Model Tutupan Lahan Di Daerah Aliran Sungai Kota Ambon Tahun 2031: Studi Kasus DAS Wai Batu Gantung, Wai Batu Gajah, Wai Tomu, Wai Batu Merah Dan Wai Ruhu'. *Jurnal Tanah dan Sumberdaya Lahan* 9(2): 473–86. <https://jtsl.ub.ac.id/index.php/jtsl/article/view/822>.
- Septory, Juan Steiven Imanuel, Philia Chisti Latue, and Heinrich Rakuasa. 2023. 'Model Dinamika Spasial Perubahan Tutupan Lahan Dan Daya Dukung Lahan Permukiman Kota Ambon Tahun 2031'. *GEOGRAPHIA: Jurnal Pendidikan dan Penelitian Geografi* 4(1): 51–62. <https://ejournal.unima.ac.id/index.php/geographia/article/view/5801>.
- Sivanpillai, Ramesh, Kevin M. Jacobs, Chloe M. Mattilio, and Ela V. Piskorski. 2021. 'Rapid Flood Inundation Mapping by Differencing Water Indices from Pre- and Post-Flood Landsat Images'. *Frontiers of Earth Science* 15(1): 1–11.

- <https://link.springer.com/10.1007/s11707-020-0818-0>.
- Soltani, Keyvan et al. 2021. 'Mapping the Spatial and Temporal Variability of Flood Susceptibility Using Remotely Sensed Normalized Difference Vegetation Index and the Forecasted Changes in the Future'. *Science of The Total Environment* 770: 145288. <https://linkinghub.elsevier.com/retrieve/pii/S0048969721003545>.
- Sugandhi, N., & Rakuasa, H. 2023. 'Utilization of Google Earth Engine for Flood Hazard Analysis in DKI Jakarta Province'. *Jurnal Riset Multidisiplin dan Inovasi Teknologi* 1(2): 40–49.
- Sugandhi, N., Rakuasa, H., Zainudin, Z., Abdul Wahab, W., Kamiludin, K., Jaelani, A., Ramdhani, R., & Rinaldi, M. 2023. 'Pemodelan Spasial Limpasan Genangan Banjir Dari DAS Ciliwung Di Kel. Kebon Baru Dan Kel. Bidara Cina DKI Jakarta'. *ULIL ALBAB : Jurnal Ilmiah Multidisiplin* 2(5): 1685–1692.
- Wulandari, Agustia Ayu, Anesia Clorita Pertuak, and Heinrich Rakuasa. 2023. 'Climate Change and Its Impact on Human Health: A Medical Geography Perspective'. *Journal of Health Science and Medical Therapy* 1(02): 80–90. <https://risetpress.com/index.php/jhsmt/article/view/279>.
- Zhao, Qiang et al. 2021. 'Progress and Trends in the Application of Google Earth and Google Earth Engine'. *Remote Sensing* 13(18): 3778. <https://www.mdpi.com/2072-4292/13/18/3778>.